

# IMMUNIZATION RECORD

Baby's Name: \_\_\_\_\_

AGE	VACCINE	DUE ON	GIVEN ON	DOCTOR'S SIGN
At Birth	BCG			
	Hepatitis B			
	Oral Polio Vaccine (OPV) – 1 <sup>st</sup> dose			
6 weeks	DPT			
	Hepatitis B			
	Hib vaccine			
	Oral Polio Vaccine (OPV) – 2 <sup>nd</sup> dose			
10 weeks	DPT			
	Hib vaccine			
	Oral Polio Vaccine (OPV) – 3 <sup>rd</sup> dose			
14 weeks	DPT			
	Hepatitis B			
	Hib vaccine			
	Oral Polio Vaccine (OPV) – 4 <sup>th</sup> dose			
6 months	Influenza vaccine – 1 <sup>st</sup> dose			
7 months	Influenza vaccine – 2 <sup>nd</sup> dose			
9 months	Measles vaccine			
>=1 year	Chickenpox vaccine			
15 months	MMR vaccine			
16–18 months	DPT – 1 <sup>st</sup> booster dose			
	Oral Polio Vaccine (OPV) – 5 <sup>th</sup> dose			
	Hib vaccine – booster dose			
2 years	Typhoid vaccine			
5 years	Typhoid vaccine			
	DPT – 2 <sup>nd</sup> booster dose			
	Oral Polio Vaccine (OPV) – 6 <sup>th</sup> dose			
10 years	TT booster dose			

BCG – Bacille Calmette-Guérin; DPT – Diphtheria, Pertussis, Tetanus; Hib – Haemophilus influenzae type b; MMR – Measles, Mumps, Rubella.

1. A dose of Hepatitis B vaccine within 12 hours of birth is essential to prevent mother-to-child transmission.
2. If Hepatitis B vaccine is given at birth, remaining doses can be given at 6 & 14 weeks, or alternatively, 6, 10 & 14 weeks (if mother is HBsAg negative).
3. Combined DPT/Hepatitis B/Hib vaccines can be given at 6, 10 & 14 weeks.
4. BCG, OPV and Hepatitis B vaccines, when missed at birth, can be started at the completion of 6 weeks.
5. For Typhoid immunization, earliest age recommended: Whole Cell vaccines at 6 months, Vi antigen at 2 years, and Oral Ty21a vaccine at 6 years. Booster dose every 3 years.
6. In addition to routine OPV doses, the recommended 'Pulse OPV dose' is also mandatory during PPI campaigns.
7. 2 doses of Influenza vaccine are given when immunizing 1st time.

## Vaccines that can be given after discussion with the parents

AGE	VACCINE	DUE ON	GIVEN ON	DOCTOR'S SIGN
>15 months	Varicella Vaccine <sup>#</sup>			
>18 months	Hepatitis A Vaccine <sup>+</sup>			
>6 weeks	Pneumococcal Conjugate Vaccine <sup>*</sup>			

<sup>#</sup><13 years of age: 1 dose; >13 years of age: 2 doses at 4–8 weeks interval. <sup>+</sup>2 doses at 6–12 months interval. <sup>\*</sup>3 primary doses at 6, 10 & 14 weeks, followed by a booster dose at 15 months.